arXiv:math/0410489v1 [math.CA] 22 Oct 2004

Potpourri, 6

Stephen William Semmes
Rice University
Houston, Texas

Let E be a nonempty finite set, and let V be a real or complex vector space. In these notes we write $\mathcal{F}(E, V)$ for the vector space of V-valued functions on E.

Suppose that E_1, E_2 are nonempty finite sets, and that V is the vector space of real or complex-valued functions on E_2. In this case we can identify $\mathcal{F}(E_1, V)$ with the vector space of real or complex-valued functions on the Cartesian product $E_1 \times E_2$, as appropriate.

In any event $\mathcal{F}(E, V)$ is basically the same as the tensor product of the vector space of real or complex-valued functions on E, as appropriate, with V. Of course the tensor product of two vector spaces, both real or both complex, is symmetric up to canonical isomorphism in the two vector spaces. With $\mathcal{F}(E, V)$ we may unravel this symmetry a bit, depending on the circumstances.

For each $x \in E$, let $\delta_x(y)$ denote the scalar-valued function on E which is equal to 1 when $y = x$ and to 0 when $y \neq x$. These functions form a basis for the scalar-valued functions on E, and in effect we may treat the vector space of scalar-valued functions on E as having a distinguished basis in this way.

We can also think of the scalar-valued functions on E as being a commutative algebra over the real or complex numbers, as appropriate, using ordinary pointwise multiplication as the product in the algebra. We can think of $\mathcal{F}(E, V)$ as being a module over the commutative algebra of scalar-valued functions, since we can multiply a vector-valued function by a scalar-valued function.

Suppose that A is any linear transformation on V. There is a unique linear transformation $\widetilde{A}$ on $\mathcal{F}(E, V)$ such that $\widetilde{A}(f)(x) = A(f(x))$ for every V-valued function f on E and every $x \in E$. Alternatively, $\widetilde{A}$ is characterized by the property that $\widetilde{A}(f\, v) = f\, A(v)$ for every scalar-valued function f on E and every vector $v \in V$.

Now suppose that T is any linear transformation acting on $\mathcal{F}(E, \mathbf{R})$ or $\mathcal{F}(E, \mathbf{C})$, as appropriate. There is a unique linear transformation $\widehat{T}$ on $\mathcal{F}(E, V)$ such that $\widehat{T}(f\, v) = T(f)\, v$ for all scalar-valued functions f on E and all $v \in V$.

Thus one can also think of $\mathcal{F}(E, V)$ as modules over the algebras of linear transformations acting on the vector space of scalar-valued functions on E and on V. The actions of $\widetilde{A}$ and $\widehat{T}$ on $\mathcal{F}(E, V)$ described in the preceding paragraphs automatically commute, by construction.

Let us focus for the moment on scalar-valued functions. Let E be a nonempty

finite set, and let $w(x)$ be a weight on E. To be more precise, $w(x)$ is a real-valued function on E such that $w(x) > 0$ for all $x \in E$.

If $f(x)$ is a real or complex-valued function on E, and p is a positive real number, put

$$\|f\|_{w,p} = \left(\sum_{x \in E} |f(x)|^p \, w(x) \right)^{1/p}. \tag{1}$$

When $w(x) = 1$ for all $x \in E$ we may simply write $\|f\|_p$ for this quantity.

For $p = \infty$ put

$$\|f\|_\infty = \|f\|_{w,\infty} = \max\{|f(x)| : x \in E\}. \tag{2}$$

Thus the weight plays no role when $p = \infty$.

Suppose that $w \equiv 1$. Clearly

$$\|f\|_\infty \le \|f\|_p \tag{3}$$

for all real or complex-valued functions f on E and all positive real numbers p. Using this one can check that

$$\|f\|_q \le \|f\|_p \tag{4}$$

when $0 < p \le q \le \infty$.

Now suppose that w defines a probability distribution on E, which is to say that

$$\sum_{x \in E} w(x) = 1. \tag{5}$$

In this event

$$\|f\|_{w,p} \le \|f\|_\infty \tag{6}$$

for all real or complex-valued functions f on E and all positive real numbers p. If r is a real number, $r \ge 1$, and $\phi(x)$ is a nonnegative real-valued function on E, then

$$\left(\sum_{x \in E} \phi(x) \, w(x) \right)^r \le \sum_{x \in E} \phi(x)^r \, w(x). \tag{7}$$

by the convexity of the function t^r on the nonnegative real numbers. It follows that

$$\|f\|_{w,p} \le \|f\|_{w,q} \tag{8}$$

when $0 < p \le q \le \infty$.

Let V be a real or complex vector space, and let $N(v)$ be a real-valued function on V such that $N(v) \ge 0$ for all $v \in V$, $N(v) = 0$ if and only if $v = 0$, and $N(\alpha \, v) = |\alpha| \, N(v)$ for all scalars α and all vectors $v \in V$. If furthermore

$$N(v + w) \le N(v) + N(w) \tag{9}$$

for all $v, w \in V$, then we say that N defines a norm on V. In the presence of the other conditions, this is equivalent to saying that the closed unit ball associated to N,

$$\{v \in V : N(v) \le 1\}, \tag{10}$$

is a convex subset of V. If there is a positive real number C such that

$$N(v+w) \le C\,(N(v) + N(w)) \tag{11}$$

for all $v, w \in V$, then we say that N defines a quasinorm on V. A special case of this occurs when there is a positive real number p, $p \le 1$, such that

$$N(v+w)^p \le N(v)^p + N(w)^p \tag{12}$$

for all $v, w \in V$.

For all p, $0 < p \le \infty$, $\|f\|_{w,p}$ satisifies the positivity and homogeneity conditions mentioned at the beginning of the previous paragraph. When $p = 1$ or $p = \infty$ one can easily verify from the definitions that $\|f\|_{w,p}$ satisifies the triangle inequality and therefore defines a norm on the vector space of real or complex-valued functions on E. The triangle inequality also holds when $1 < p < \infty$, as one can check by showing that the associated closed unit ball is convex using the convexity of t^p on the nonnegative real numbers. When $0 < p \le 1$ one has that

$$\|f_1 + f_2\|_{w,p}^p \le \|f_1\|_{w,p}^p + \|f_2\|_{w,p}^p \tag{13}$$

for all real or complex-valued functions f_1, f_2 on E. This follows from the fact that

$$(a+b)^p \le a^p + b^p \tag{14}$$

for all nonnegative real numbers a, b.

Suppose that f_1, f_2 are real or complex-valued functions on E, $1 \le p, q \le \infty$, and

$$\frac{1}{p} + \frac{1}{q} = 1. \tag{15}$$

If either p or q is infinite, then its reciprocal is equal to 0. Hölder's inequality states that

$$\Big| \sum_{x \in E} f_1(x)\, f_2(x)\, w(x) \Big| \le \|f_1\|_{w,p}\, \|f_2\|_{w,q} \tag{16}$$

in this case. Of course one might as well restrict one's attention to functions f_1, f_2 which are real-valued and nonnegative. If $p = 1$ and $q = \infty$, or $q = 1$ and $p = \infty$, then Hölder's inequality can be verified directly.

Suppose that $1 < p, q < \infty$. If a, b are nonnegative real numbers, then

$$a\,b \le \frac{a^p}{p} + \frac{b^q}{q}. \tag{17}$$

This can be seen as a special case of the convexity of the exponential function, for instance. Hence

$$|f_1(x)\, f_2(x)| \le \frac{|f_1(x)|^p}{p} + \frac{|f_2(x)|^q}{q} \tag{18}$$

for all $x \in E$, and therefore

$$\Big|\sum_{x \in E} f_1(x)\, f_2(x)\, w(x)\Big| \le \frac{\|f_1\|_{w,p}^p}{p} + \frac{\|f_2\|_{w,q}^q}{q}. \tag{19}$$

This implies Hölder's inequality when $\|f_1\|_{w,p}$ and $\|f_2\|_{w,q}$ are equal to 1, and the general case follows from a scaling argument.

If f_1, f_2 are functions on E and $0 < p, q, r \le \infty$,

$$\frac{1}{r} = \frac{1}{p} + \frac{1}{q}, \tag{20}$$

then

$$\|f_1\, f_2\|_{w,r} \le \|f_1\|_{w,p}\, \|f_2\|_{w,q}, \tag{21}$$

by Hölder's inequality, with the obvious conventions for infinite exponents. For a single function f on E we have that

$$\|f\|_{w,r} \le \|f\|_{w,p}^{p/r}\, \|f\|_{w,q}^{q/r}, \tag{22}$$

again by Hölder's inequality. One can also use Hölder's inequality to check that $\|f\|_{w,p}$ satisfies the triangle inequality when $1 < p < \infty$, also known as Minkowski's inequality. Namely,

$$\begin{aligned} &\sum_{x \in E} |f_1(x) + f_2(x)|^p\, w(x) \\ &\le \sum_{x \in E} |f_1(x)|\, |f_1(x) + f_2(x)|^{p-1}\, w(x) + \sum_{x \in E} |f_2(x)|\, |f_1(x) + f_2(x)|^{p-1}\, w(x), \end{aligned} \tag{23}$$

and one can apply Hölder's inequality to each of the sums on the right side.

Let h be a real or complex-valued function on E, and define a linear transformation M_h on the vector space of real or complex-valued functions on E, as appropriate, by

$$M_h(f)(x) = h(x)\, f(x), \tag{24}$$

so that $M_h(f)$ is the product of h and f. If p, q, r are as in the preceding paragraph, then we get that

$$\|M_h(f)\|_{w,r} \le \|h\|_{w,q}\, \|f\|_{w,p} \tag{25}$$

for all functions f on E. This inequality is sharp in the sense that for each nonzero function h on E there is a nonzero function f on E such that equality holds and indeed such that

$$|h(x)\, f(x)|^r = |h(x)|^q = |f(x)|^p \tag{26}$$

for all $x \in E$, at least when $p, q, r < \infty$. For infinite exponents one can make simple adjustments to this last part.

Let T be a linear transformation on the vector space of real or complex-valued functions on E, and fix r, s with $0 < r, s \leq \infty$. Suppose that k is a nonnegative real number such that

$$\|T(f)\|_{w,s} \leq k \, \|f\|_{w,r} \tag{27}$$

for all functions f on E. Let h_1, h_2 be real or complex-valued functions on E, as appropriate, and suppose that $0 < q_1, q_2 \leq \infty$ and that $q_1 \geq r$. Choose p, t with $0 < p, t \leq \infty$ such that

$$\frac{1}{r} = \frac{1}{p} + \frac{1}{q_1} \tag{28}$$

and

$$\frac{1}{t} = \frac{1}{q_2} + \frac{1}{s}. \tag{29}$$

It follows from the earlier inequalities that

$$\|(M_{h_2} \circ T \circ M_{h_1})(\phi)\|_{w,t} \leq k \, \|h_1\|_{w,q_1} \, \|h_2\|_{w,q_2} \, \|\phi\|_{w,p} \tag{30}$$

for all functions ϕ on E. Conversely, suppose that k is a nonnegative real number such that this inequality holds for all functions h_1, h_2, and ϕ on E. Then one can check that $\|T(f)\|_{w,s} \leq k \, \|f\|_{w,r}$ for all functions f on E. This is basically the same as the sharpness of the inequality for the norm of $M_h(f)$ discussed in the previous paragraph.

Now suppose that V is a real or complex vector space with quasinorm norm $\| \cdot \|_V$. If $0 < p \leq \infty$ and f is a V-valued function on E, put

$$\|f\|_{w,p,V} = \Big(\sum_{x \in E} \|f(x)\|_V^p \, w(x) \Big)^{1/p} \tag{31}$$

when $p < \infty$ and

$$\|f\|_{\infty,V} = \|f\|_{w,\infty,V} = \max\{\|f(x)\|_V : x \in E\}. \tag{32}$$

This is the same as $\| \cdot \|_{w,p}$ applied to the real-valued function $\|f(x)\|_V$ on E.

One can check that $\|f\|_{w,p,V}$ is a quasinorm on $\mathcal{F}(E,V)$ for all p. If $\|v\|_V$ is a norm on V and $1 \leq p \leq \infty$, then $\|f\|_{w,p,V}$ is a norm on $\mathcal{F}(E,V)$. If $0 < p \leq 1$ and $\|v\|_V$ satisfies

$$\|v + w\|_V^p \leq \|v\|_V^p + \|w\|_V^p \tag{33}$$

for all $v, w \in V$, then $\|f\|_{w,p,V}$ satisfies the analogous condition on $\mathcal{F}(E,V)$. Observe that if $\|v\|_V$ satisfies this condition and $0 < q \leq p$, then we also have that

$$\|v + w\|_V^q \leq \|v\|_V^q + \|w\|_V^q \tag{34}$$

for all $v, w \in V$.

Let h be a real or complex-valued function on E, as appropriate, and let us again write M_h for the linear operator on $\mathcal{F}(E, V)$ given by multiplication by h. Thus $M_h(f)(x) = h(x)\, f(x)$ for all V-valued functions f on E. If $0 < p, q, r \le \infty$ and $1/r = 1/p + 1/q$, then

$$\|M_h(f)\|_{w,r,V} \le \|h\|_{w,q}\, \|f\|_{w,p,V} \tag{35}$$

for all V-valued functions f on E.

As in the scalar case, suppose that $0 < q_1, q_2, r, s \le \infty$ and $q_1 \ge r$, and choose p, t with $0 < p, t \le \infty$ so that $1/r = 1/p + 1/q_1$ and $1/t = 1/q_2 + 1/s$. If Z is a linear transformation on $\mathcal{F}(E, V)$ and k is a nonnegative real number, then

$$\|Z(f)\|_{w,s,V} \le k\, \|f\|_{w,r,V} \tag{36}$$

for all V-valued functions f on E if and only if

$$\|(M_{h_2} \circ Z \circ M_{h_1})(\phi)\|_{w,t,Z} \le k\, \|h_1\|_{w,q_1}\, \|h_2\|_{w,q_2}\, \|\phi\|_{w,s,V} \tag{37}$$

for all real or complex-valued functions h_1, h_2 on E, as appropriate, and all V-valued functions ϕ on E. In particular this applies to the case where $Z = \widehat{T}$ for some linear transformation T acting on the vector space of real or complex-valued functions on E, as appropriate. In this event $M_{h_2} \circ Z \circ M_{h_1}$, as a linear transformation on V-valued functions, is the same as

$$\widehat{R},\ R = M_{h_2} \circ T \circ M_{h_1}, \tag{38}$$

where M_{h_1}, M_{h_2} refer to the multiplication operators on scalar-valued functions in the definition of R.

Let $a(x, y)$ be a real or complex-valued function on $E \times E$. This leads to a linear transformation A acting on real or complex-valued functions on E, as appropriate, given by

$$A(f)(x) = \sum_{y \in E} a(x, y)\, f(y)\, w(y), \tag{39}$$

and every such linear transformation can be expressed in this manner. Consider the condition

$$\|A(f)\|_{w,1} \le \|f\|_{w,\infty} \tag{40}$$

for all functions f on E. This holds in particular if

$$\sum_{x \in E} \sum_{y \in E} |a(x, y)|\, w(x)\, w(y) \le 1. \tag{41}$$

Suppose that T is a linear transformation acting on real or complex-valued functions on E. Let r, s be exponents with $1 \le r, s \le \infty$, and assume that

$$\|T(f)\|_{w,s} \le \|f\|_{w,r} \tag{42}$$

for all functions f on E. Let q, $1 \le q \le \infty$, be the exponent conjugate to s, so that $1/q + 1/s = 1$, and let h_1, h_2 be real or complex-valued functions on E, as appropriate, such that

$$\|h_1\|_{w,r}, \|h_2\|_{w,q} \le 1. \tag{43}$$

Under these conditions, the linear transformation

$$A = M_{h_2} \circ T \circ M_{h_1} \tag{44}$$

acting on functions on E satisfies the conditions described in the previous paragraph.

Let W be a finite-dimensional real or complex vector space equipped with a norm $\|\cdot\|_W$. Consider expressions for linear transformations on W of the form

$$\sum_{\ell=1}^{n} c_\ell \, \lambda_\ell \, w_\ell, \tag{45}$$

where the coefficients c_ℓ are real or complex numbers, as appropriate, the w_ℓ's are elements of W with $\|w_\ell\|_W \le 1$ for all ℓ, and the λ_ℓ's are linear functionals on W such that $|\lambda_\ell(u)| \le \|u\|_W$ for all $u \in W$, which is to say that the λ_ℓ's have dual norm less than or equal to 1. Every linear transformation on W can be expressed as a sum of this kind.

By definition, the trace norm of a linear transformation on W is equal to the infimum of

$$\sum_{\ell=1}^{n} |\alpha_\ell| \tag{46}$$

over all representations of the linear transformation as in the previous paragraph. Let p be a positive real number such that $p \le 1$. We can consider more generally the infimum of

$$\Big(\sum_{\ell=1}^{n} |\alpha_\ell|^p\Big)^{1/p} \tag{47}$$

over all such representations of a given linear transformation. This quantity is greater than or equal to the trace norm, and is monotonically increasing as p decreases. It defines a quasinorm on the vector space of linear transformations on W which satisfies the generalization of the triangle inequality in which the pth power of the quasinorm is subadditive.

References

[Bea1] R. Beals, *Topics in Operator Theory*, The University of Chicago Press, 1971.

[Bea2] R. Beals, *Advanced Mathematical Analysis: Periodic Functions and Distributions, Complex Analysis, Laplace Transform and Applications*, Springer-Verlag, 1973.

[BenS] C. Bennett and R. Sharpley, *Interpolation of Operators*, Academic Press, 1988.

[BenL] Y. Benyamini and J. Lindenstrauss, *Geometric Nonlinear Functional Analysis*, Volume 1, Colloquium Publications **48**, American Mathematical Society, 2000.

[BerL] J. Bergh and J. Löfström, *Interpolation Spaces*, Springer-Verlag, 1976.

[Boc] S. Bochner, *Lectures on Fourier Integrals*, with an author's supplement on monotonic functions, Stieltjes integrals, and harmonic analysis, translated from the 1932 original by M. Tenenbaum and H. Pollard, Annals of Mathematics Studies **42**, Princeton University Press, 1959.

[BocC] S. Bochner and K. Chandrasekharan, *Fourier Transforms*, Annals of Mathematics Studies **19**, Princeton University Press, 1949.

[Duo] J. Duoandikoetxea, *Fourier Analysis*, translated and revised by D. Cruz-Uribe, SFO, American Mathematical Society, 2001.

[GarR] J. García-Cuerva and J. Rubio de Francia, *Weighted Norm Inequalities and Related Topics*, North-Holland, 1985.

[Gol] R. Goldberg, *Methods of Real Analysis*, 2nd edition, Wiley, 1976.

[HewS] E. Hewitt and K. Stromberg, *Real and Abstract Analysis*, Springer-Verlag, 1975.

[JohL] W. Johnson and J. Lindenstrauss, editors, *Handbook of the Geometry of Banach Spaces*, Volumes I and II, North-Holland, 2001 and 2003.

[Jon] F. Jones, *Lebesgue Integration on Euclidean Space*, Jones and Bartlett, 1993.

[Jou] J.-L. Journé, *Calderón–Zygmund Operators, Pseudodifferential Operators, and the Cauchy Integral of Calderón*, Lecture Notes in Mathematics **994**, Springer-Verlag, 1993.

[KalPR] N. Kalton, N. Peck, and J. Roberts, *An F-Space Sampler*, Cambridge University Press, 1984.

[Kat] Y. Katznelson, *An Introduction to Harmonic Analysis*, 3rd edition, Cambridge University Press, 2004.

[Kra1] S. Krantz, *Real Analysis and Foundations*, CRC Press, 1991.

[Kra2] S. Krantz, *A Panorama of Harmonic Analysis*, Carus Mathematical Monographs **27**, Mathematical Association of America, 1999.

[LinT] J. Lindenstrauss and L. Tzafriri, *Classical Banach Spaces I, Sequence Spaces, II, Function Spaces*, Springer-Verlag, 1977, 1979.

[McC] C. McCarthy, c_p, Israel Journal of Mathematics **5** (1967), 249–271.

[Mil] M. Milman, *Extrapolation and Optimal Decompositions with Applications to Analysis*, Lecture Notes in Mathematics **1580**, Springer-Verlag, 1994.

[MilS] V. Milman and G. Schechtman, *Asymptotic Theory of Finite-Dimensional Normed Spaces*, with an appendix by M. Gromov, Lecture Notes in Mathematics **1200**, Springer-Verlag, 1986.

[Pelc] A. Pełczyński, *Banach Spaces of Analytic Functions and Absolutely Summing Operators*, Conference Board of the Mathematical Sciences Regional Conference Series in Mathematics **30**, American Mathematical Society, 1977.

[Pell] V. Peller, *Hankel Operators and their Applications*, Springer-Verlag, 2003.

[Pis1] G. Pisier, *Factorization of Linear Operators and Geometry of Banach Spaces*, Conference Board of the Mathematical Sciences Regional Conference Series in Mathematics **60**, American Mathematical Society, 1986.

[Pis2] G. Pisier, *The Volume of Convex Bodies and Banach Space Geometry*, Cambridge University Press, 1989.

[Rud1] W. Rudin, *Principles of Mathematical Analysis*, 3rd edition, McGraw-Hill, 1976.

[Rud2] W. Rudin, *Real and Complex Analysis*, 3rd edition, McGraw-Hill, 1987.

[Rud3] W. Rudin, *Functional Analysis*, 2nd edition, McGraw-Hill, 1991.

[Sad] C. Sadosky, *Interpolation of Operators and Singular Integrals: An Introduction to Harmonic Analysis*, Marcel Dekker, 1979.

[Sim] B. Simon, *Trace Ideals and their Applications*, Cambrdige University Press, 1979.

[Ste1] E. Stein, *Singular Integrals and Differentiability Properties of Functions*, Princeton University Press, 1970.

[Ste2] E. Stein, *The development of square functions in the work of A. Zygmund*, Bulletin of the American Mathematical Society (New Series) **7** (1982), 359–376.

[Ste3] E. Stein, *Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals*, Princeton University Press, 1993.

[SteS1] E. Stein and R. Shakarchi, *Fourier Analysis: An Introduction*, Princeton University Press, 2003.

[SteS2] E. Stein and R. Shakarchi, *Complex Analysis*, Princeton University Press, 2003.

[SteS3] E. Stein and R. Shakarchi, *Real Analysis*, Princeton University Press, to appear.

[SteS4] E. Stein and R. Shakarchi, *Probability Theory*, Princeton University Press, to appear.

[SteW1] E. Stein and G. Weiss, *Interpolation of operators with change of measures*, Transactions of the American Mathematical Society **87** (1958), 159–172.

[SteW2] E. Stein and G. Weiss, *Introduction to Fourier Analysis on Euclidean Spaces*, Princeton University Press, 1971.

[StrT] J.-O. Strömberg and A. Torchinsky, *Weighted Hardy Spaces*, Lecture Notes in Mathematics **1381**, Springer-Verlag, 1989.

[Tor] A. Torchinsky, *Real-Variable Methods in Harmonic Analysis*, Dover, 2004.

[Tre] F. Treves, *Topological Vector Spaces, Distributions, and Kernels*, Academic Press, 1967.

[WheZ] R. Wheeden and A. Zygmund, *Measure and Integral: An Introduction to Real Analysis*, Marcel Dekker, 1977.

[Woj] P. Wojtaszczyk, *Banach Spaces for Analysts*, Cambridge University Press, 1991.

[Zem] A. Zemanian, *Distribution Theory and Transform Analysis: An Introduction to Generalized Functions, with Applications*, 2nd edition, Dover, 1987.

[Zyg] A. Zygmund, *Trigonometric Series*, Volumes I and II, 3rd edition, Cambridge University Press, 2002.